

11:20 – 11:50

S6: POSTERS SESSION

11:20 – 11:27 S6-1 (PP)

SHOULD UPPER LIP GRAFT (ULG) BE THE FIRST OPTION FOR REDO HYPOSPADIAS IN CHILDREN?

Tariq Osman ABBAS¹, A ELKADHI, Bruno LESLIE, Santiago VALLASCIANI and Joao PIPPI SALLE

Sidra Medicine, Urology, Doha, QATAR

11:27 – 11:34 S6-2 (PP)

POSTERIOR URETHRAL VALVES IN PATIENTS WITH TRISOMY 21: A FREQUENT ASSOCIATION WITH DIVERSE PRESENTATION

Harriet CORBETT¹, Charlotte KANE², Anna-May LONG² and Helen Fiona MCANDREW¹

1 Alder Hey Children's NHS Foundation Trust, Regional Department of Urology, Liverpool, UNITED KINGDOM

2 Alder Hey Children's NHS Foundation Trust, Department of Surgery, Liverpool, UNITED KINGDOM

11:34 – 11:41 S6-3 (PP)

EFFECTS OF A 2-STEP MOBILE APPLICATION PROGRAM ON SELF-MANAGEMENT AND QUALITY OF LIFE OF CHILDREN WITH SPINA BIFIDA

Eun Kyoung CHOI¹, Eunjeong BAE², Eunyong JUNG², Yong Ju LEE³, Yoonhye JI⁴ and Sang Won HAN⁵

1 Yonsei University, College of Nursing, Seoul, REPUBLIC OF KOREA

2 Yonsei University Graduate School, Department of Nursing, Seoul, REPUBLIC OF KOREA

3 Hallym University, Kangnam Sacred Heart Hospital, Department of Pediatrics, Seoul, REPUBLIC OF KOREA

4 Severance Children's Hospital, Yonsei University Health System, Department of Pediatric Urology, Seoul, REPUBLIC OF KOREA

5 Yonsei University College of Medicine, Department of Urology and Urological Science Institute, Seoul, REPUBLIC OF KOREA

11:41 – 11:48 S6-4 (PP)

A LOWER D2/D4 DIGIT RATIO AND SCROTAL BASE DISTANCE IN MALE INFANTS WITH HYPOSPADIAS AND/OR CRYPTORCHIDISM THAN NORMAL BOYS

Tariq Osman ABBAS¹, A ELKADHI² and Joao PIPPI SALLE³

1 Hamad General Hospital, Pediatric Surgery, Doha, QATAR

2 Hamad Medical Corporation, Pediatric Surgery, Doha, QATAR

3 Sidra Medicine, Urology, Doha, QATAR